

The failure of cosmological perturbation theory in the new ekpyrotic and cyclic ekpyrotic scenarios

David H. Lyth

Physics Department, Lancaster University, Lancaster LA1 4YB, U.K.

Abstract

Cosmological perturbation theory fails in the new ekpyrotic and cyclic ekpyrotic scenarios, before the scale factor of the Universe reaches zero. As a result, a recently-proposed recipe for evolving the curvature perturbation through the bounce in these scenarios cannot be justified.

Introduction It is now clear that the dominant cause of large-scale structure in the Universe is a perturbation in the spatial curvature [1]. The perturbation exists already when cosmological scales are approaching horizon entry, at which stage it has an almost flat spectrum. It presumably originates during an early era of superluminal evolution, defined as one in which comoving scales are leaving the horizon, at the beginning of which the entire observable Universe is inside the horizon. During superluminal evolution the curvature of the Universe in Hubble units is decreasing, explaining naturally the near-flatness of the observable Universe.

The usual hypothesis concerning the superluminal era is that it is one of expansion, or in other words an era of inflation, with slowly varying Hubble parameter H . During inflation, each canonically-normalized light scalar field acquires a perturbation with a flat spectrum, and this provides a natural origin for the observed curvature perturbation. The curvature perturbation might be generated directly during inflation from the perturbation in the slowly-rolling inflaton field [1], or indirectly [2] by a different field (curvaton) whose isocurvature density perturbation is converted into a curvature perturbation during some radiation-dominated era after inflation.

The alternative to inflation is an era of superluminal collapse (negative Hubble parameter) giving way at some stage to expansion (positive Hubble parameter). Unless spatial curvature becomes significant near the bounce, the theory describing the bounce cannot be standard field theory with Einstein gravity, corresponding to an action of the form

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} M_{\text{P}}^2 R - \frac{1}{2} g^{\mu\nu} G_{IJ}(\phi^K) \partial_\mu \phi^I \partial_\nu \phi^J - V(\phi^K) \right]. \quad (1)$$

The reason is that the energy density ρ and pressure P in such a theory satisfies the null energy condition $\rho + P > 0$, which implies that the energy density decreases monotonically. Through the Friedmann equation, this implies that the Hubble parameter decreases monotonically in the absence of spatial curvature, forbidding any bounce. The bounce might be described by a non-standard four-dimensional field theory, but more plausibly would involve extra dimensions and/or string theory.

The pre-big-bang scenario Until recently, only one type of superluminal collapse has been considered, and the scenario involving it is termed simply the pre-big-bang scenario [3, 4]. In this scenario the action is supposed to have the standard form Eq. (1) during both collapse and expansion, up to some maximum energy density ρ_b . This is the bounce energy density, which is supposed to correspond to the string scale and is at or below the Planck scale $\rho_P = M_P^4 = (2 \times 10^{18} \text{ GeV})^4$. During pre-big-bang collapse, the energy density of the Universe comes from a scalar field ϕ with no potential. Taking time t as negative this leads to scale factor $a \propto |t|^{\frac{1}{3}}$ corresponding to $H \equiv \dot{a}/a = 1/3t$. The same result is obtained if there are several such fields, each giving a fixed fraction of the energy density.

The ekpyrotic scenario Recently a different mechanism for superluminal collapse has been proposed, termed the ekpyrotic mechanism [5, 6]. It invokes an extra dimension in which two branes are attracted towards each other. After integrating out the extra dimension one obtains superluminal four-dimensional collapse, in which the energy density and pressure are dominated by a canonically-normalized field ϕ which corresponds to the distance between the branes. The action is of the standard form Eq. (1),

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} M_P^2 R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) + \dots \right], \quad (2)$$

the remaining terms having negligible effect on the evolution of ϕ .

From some very early time, until after cosmological scales have left the horizon, the potential during collapse is taken to be

$$V \simeq -V_0 \exp \left(-\sqrt{\frac{2}{p}} \frac{\phi}{M_P} \right) \quad (3)$$

with $p \ll 1$. It is supposed that $\dot{\phi}$ starts out from a sufficiently small value, leading to power-law collapse $a \propto |t|^p$ in which the Hubble parameter and energy density are related by the Friedmann equation

$$3M_P^2 H^2 = \rho \quad (4)$$

$$= \frac{1}{2} \dot{\phi}^2 + V(\phi). \quad (5)$$

The energy density increases during collapse, but it is always small compared with $|V|$ by virtue of the condition $p \ll 1$.

Three versions of the ekpyrotic scenario have been proposed, in references [5], [6] and [7], which we refer to as respectively the old, new and cyclic ekpyrotic scenarios. In each of them it is claimed that the observed curvature perturbation can be generated by the quantum fluctuation of the field ϕ . In an earlier note [8] we demonstrated that this claim is invalid in the case of the old ekpyrotic scenario, as described in [5]. The purpose of the present note is to show that the claim is also invalid in the case of the new ekpyrotic scenario as described in reference [6], because in that reference linear cosmological perturbation theory is being invoked in a regime where it is not valid. Since the cyclic ekpyrotic Universe [7] invokes the same mechanism for producing the density perturbation, the same statement applies to that scenario also.

The comoving curvature perturbation To define the cosmological perturbations one has to choose a gauge (coordinate system) which defines a threading and slicing of spacetime. Let us denote a generic perturbation by the symbol g . At each moment of time, it is Fourier-expanded in comoving coordinates

$$g(\mathbf{x}) = (2\pi)^{-\frac{3}{2}} \int g(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}} d^3\mathbf{k}, \quad (6)$$

with k/a the wavenumber. Its spectrum is defined by [9, 10, 1]

$$\langle g(\mathbf{k}) g^*(\mathbf{k}') \rangle = 2\pi^2 k^{-3} \delta^3(\mathbf{k} - \mathbf{k}') \mathcal{P}_g(k), \quad (7)$$

where the brackets denote an ensemble average. (On the usual supposition that perturbations originate as quantum fluctuations, the brackets denote the quantum expectation value.) The formal expectation value of $g^2(\mathbf{x})$ at any point in space is

$$\langle g^2(\mathbf{x}) \rangle = \int_0^\infty \mathcal{P}_g(k) d(\ln k), \quad (8)$$

and $\mathcal{P}_g^{\frac{1}{2}}$ is the typical magnitude of a fluctuation in $g(\mathbf{x})$ with size of order $1/k$. The perturbation has a time-dependence, which is inherited by the spectrum,

$$\mathcal{P}_g^{\frac{1}{2}}(\mathbf{k}, t) = \text{const} |g(\mathbf{k}, t)|. \quad (9)$$

To define the spatial curvature perturbation we need a slicing of spacetime, which is best taken as the one orthogonal to comoving worldlines, the comoving slicing. Its line element may be written

$$d\ell^2 = a^2(t)(1 + 2\mathcal{R})\delta_{ij}dx^i dx^j, \quad (10)$$

and $\mathcal{R}$ defines the curvature perturbation.¹ This is the best choice because under very general circumstances $\mathcal{R}$ is time-independent on super-horizon scales. As a result, the

¹This is the ϕ_m of [11]. On the super-horizon scales of interest the comoving slicing is practically coincident with the slicing of uniform energy density which can equally well be used. The corresponding perturbation is the ζ of [12].

curvature perturbation that is generated in the very early Universe should be the one that is observed as cosmological scales start to enter the horizon.

There is also a slicing with no curvature perturbation, the flat slicing. If the displacement from the flat slicing to the comoving one is $\delta t(\mathbf{x}, t)$ the curvature perturbation on the latter is $\mathcal{R} = |H|\delta t$. (A similar formula actually gives the curvature perturbation on any slicing.) Whenever the energy density and pressure are dominated by single field ϕ , this field is uniform on comoving slices [12]. It follows [13, 15] that $\mathcal{R} = -|H|\delta\phi/\dot{\phi}$, where $\delta\phi$ is defined on the spatially flat slicing. (It is the ‘gauge-invariant’ field perturbation of [15].)

The curvature perturbation in the pre-big-bang scenario The spectrum of the curvature perturbation during superluminal evolution is found by a standard method [1]. In the pre-big-bang case one finds [3, 4], well after the epoch $aH = k$ of horizon exit,

$$\mathcal{P}_{\mathcal{R}}^{\frac{1}{2}} = \frac{1}{2\sqrt{6}\pi^{\frac{3}{2}}} \left(\frac{|H|}{M_{\text{P}}} \right) \left(\frac{k}{a|H|} \right)^{\frac{3}{2}} \ln \left(\frac{a|H|}{k} \right). \quad (11)$$

This spectrum is time-independent except for the logarithm, and corresponds to spectral index $n = 4$.

Assuming that the curvature perturbation is continuous across the bounce it has to be generated in the pre-big-bang scenario after the bounce by some curvaton field. To possess a flat spectrum in the face of the rapidly-varying H , this field must have [16, 4, 17, 2] a suitable initial condition, and a suitable coupling to ϕ . An alternative proposal might be to generate the curvature perturbation (or conceivably a curvaton field perturbation) during the bounce, but such a thing has never been suggested let alone a mechanism for achieving it.

The curvature perturbation in the old ekpyrotic scenario In the old ekpyrotic scenario [5], it is supposed that the brane collision corresponds to some finite field value ϕ , corresponding to energy density ρ . After the collision the Universe is expanding with the same initial energy density and there is no ϕ field. In this scenario the spectrum of the curvature perturbation during collapse can be calculated in exactly the same way as for the pre-big-bang, but taking into account the exponential potential.² Well after horizon exit, to leading order in k/aH , the spectrum [8] is time-independent with spectral index $n \simeq 3$. This result is specific to the exponential potential Eq. (3) but there is no reason to think that a different potential would give the necessary flat spectrum and in any case something like Eq. (3) is motivated by string theory. The conclusion therefore is that in the old ekpyrotic scenario, the observed curvature perturbation has to be generated either at the bounce or subsequently by a curvaton field.

This conclusion is different from the one reached in the original ekpyrotic paper [5] (see also [18]). That paper ignored the effect of the metric perturbation on the field evolution

²In [5] it is stated that just before collision the potential climbs abruptly back to zero, but this has a negligible effect on the result.

(gravitational back-reaction), and arrived at the incorrect conclusion that the comoving curvature perturbation generated during collapse has a flat spectrum, so that it could be the observed curvature perturbation.³

The new and cyclic ekpyrotic scenarios In the new [6] ekpyrotic scenario it is supposed, on the basis of string theory, that the brane collision corresponds to $\phi = -\infty$. There are two stages of collapse. In the first stage, which ends only after cosmological scales have left the horizon, the potential is given by Eq. (3) as in the old ekpyrotic scenario. In the second stage, the potential climbs towards zero and becomes negligible compared with the kinetic term so that the evolution of perturbations is the same as in the pre-big-bang scenario. This stage ends when $\phi = -\infty$, corresponding to infinite energy density, infinite Hubble parameter and zero scale factor a . The branes then collide and the expansion of the Universe begins, but in contrast with the old ekpyrotic scenario the branes move apart after the collision so that the field ϕ continues to exist, though with a possibly different potential. The potential is at least initially negligible so that the contribution of ϕ to the energy density falls like a^{-6} . In addition there is the energy density of radiation produced by the brane collision, which falls like a^{-4} and eventually dominates leading to the usual Hot Big Bang.

The cyclic ekpyrotic scenario [7] is the same, except that the potential of ϕ is the same during expansion as during collapse, and this potential asymptotes at late times to a nonzero positive constant. The latter feature gives rise to a slowly-increasing cosmological constant and eventually to the re-collapse of the Universe and therefore to a continuous cycle of expansion and collapse. None of this matters for the present purpose, because the mechanism for generating the observed curvature perturbation proposed in [7] is the same as the one proposed in [6]. Accordingly we refer from now on to the ‘new’ ekpyrotic scenario, taking it as read that the statements apply also to the cyclic ekpyrotic scenario.

The general idea of the new ekpyrotic scenario, developed in an earlier related paper [21], is that the underlying five-dimensional theory is non-singular as one goes through the bounce, so that it may in some sense be permissible to allow the four-dimensional scale factor to go through zero. The standard field theory action Eq. (1) is supposed in some sense to remain valid all the way down to $a = 0$. It is of course well-known that Eq. (1) cannot be literally valid on time- and distance scales below the Planck scale, because the quantum fluctuation on such scales is out of control. For instance, on the distance scale M_{P}^{-1} the vacuum fluctuation in the energy density corresponding to the field gradients is of order M_{P}^4 , which is big enough in relation to the distance scale to form black holes (the spacetime foam). In particular, the theory cannot be taken literally when the Hubble time falls below the Planck scale, corresponding to $H \gtrsim M_{\text{P}}$ and $\rho \gtrsim M_{\text{P}}^4$.

³The words ‘curvature perturbation’ do not appear in [5], which uses the formalism of Guth and Pi [19] and Olson [20]. However, as pointed out in [8], that formalism is equivalent to the usual one, because of the formula $\mathcal{R} = |H|\delta t$ where δt is the time delay introduced by Guth and Pi. We emphasize that the equivalence is complete provided that $\mathcal{R}$ is correctly evaluated. The approximation that δt is constant, made by Guth and Pi, is inessential though it is valid during inflation in the slow-roll approximation. Usually, $\mathcal{R}$ is constant on super-horizon scales, which requires $\delta t \propto 1/H$.

This problem is clearly recognized in [21] (“there is concern that the quantum fluctuations become uncontrolled at $t = 0$ ”).

Nevertheless, the action of Eq. (2) is used in [6] to calculate a curvature perturbation which, after the bounce, has the flat spectrum required by observation. The purpose of this note is to point out that the calculation is invalid, because it invokes linear cosmological perturbation theory outside its regime of validity.

The calculation of reference [6] The calculation in [6] starts with the Bardeen potential [11], which is usually defined as the spatial curvature perturbation in the conformal Newtonian gauge;

$$ds^2 = a^2(t) \left(-(1 + 2\Psi)d\tau^2 + (1 - 2\Phi)\delta_{ij}dx^i dx^j \right). \quad (12)$$

Alternatively, it may be defined in terms of the density contrast $\delta \equiv \delta\rho/\rho$ defined on comoving slices (the Bardeen [11] variable ϵ_m),

$$\delta = -\frac{2}{3} \left(\frac{k}{aH} \right)^2 \Phi. \quad (13)$$

The definitions are equivalent within linear perturbation theory, but the first one requires $|\Phi| \ll 1$ which is quite restrictive [11] and the second one is preferable.

The spectrum of Φ during the first stage of collapse may in principle be calculated in exactly the same way as the spectrum of $\mathcal{R}$, using the evolution equation for the field perturbation $\delta\phi_N$ and imposing the same condition of an initial vacuum state. (A simpler technique [22, 6] is to consider the evolution of Φ , using its relation with $\delta\phi_N$ only to impose the initial vacuum condition. A third method would be to deduce $\mathcal{P}_\Phi$ from the (slowly-varying) $\mathcal{P}_\mathcal{R}$, using Eq. (9) with Eq. (17) below.) It turns out that during this stage the evolution of $\delta\phi_N$ is not much affected by gravitational back-reaction and one finds

$$\mathcal{P}_\Phi^{\frac{1}{2}} = \frac{1}{2\pi} \sqrt{\frac{1}{2p}} \frac{H}{M_P} \left(\frac{aH}{k} \right)^p. \quad (14)$$

The last factor gives the effect of gravitational back-reaction.

The idea behind the proposal of [6] is that the flat spectrum Eq. (14) of the Bardeen potential may be converted at the bounce into a flat spectrum for $\mathcal{R}$, through the coupled evolution equations of Einstein gravity. In these equations, anisotropic stress may presumably be ignored on the super-horizon scales of interest. The coupled equations are then [9, 10, 1]

$$-\mathcal{R} = \frac{2}{3} \frac{|H|^{-1} \dot{\Phi} + \Phi}{1 + w} + \Phi \quad (15)$$

$$|H|^{-1} \dot{\mathcal{R}} = -\frac{\delta P}{\rho + P}, \quad (16)$$

where δP is defined on the comoving slicing. The non-adiabatic pressure perturbation $\delta P_{\text{nad}} \equiv \delta P - (\dot{P}/\rho)\delta\rho$ is also ignored on on super-horizon scales, which is appropriate in

a scenario where one is trying to generate the curvature perturbation without a curvaton [2, 17] field. Then Eq. (16) becomes

$$|H|^{-1}\dot{\mathcal{R}} = -\frac{c_s^2}{1+w}\delta \equiv \frac{3}{2}\frac{c_s^2}{1+w}\left(\frac{k}{aH}\right)^2\Phi, \quad (17)$$

where $c_s^2 \equiv \dot{P}/\dot{\rho}$.

Eqs. (15) and (17) form a closed system with two independent solutions. From Eq. (17) it follows that $\mathcal{R}$ is *practically constant provided that* $c_s^2|\delta| \ll (1+w)|\mathcal{R}|$. This condition is very well satisfied in the expanding Universe for any reasonable initial condition and any reasonable behavior of a and w [10, 1]. It is satisfied during the first stage of ekpyrotic collapse and marginally violated during pre-big-bang collapse.

During both stages of ekpyrotic collapse the left hand side of Eq. (15) is negligible leading to $\Phi \propto H/a$. Using this result and Eqs. (9) and (14), the spectrum of δ in the limit $t \rightarrow 0$ when the potential has become negligible is given by $\mathcal{P}_\delta^{\frac{1}{2}} \rightarrow Ak^{1+p}$ with A a constant. Using Eqs. (9) and (15) this implies that the spectrum of $\mathcal{R}$ is logarithmically divergent, $6\mathcal{P}_\mathcal{R}^{\frac{1}{2}} \rightarrow \mathcal{P}_\delta^{\frac{1}{2}} \ln|t|$.

Now we come to the crucial point. Because $\mathcal{R}$ is divergent, the authors of [6] focus on δ . Combining Eqs. (17) and (15) gives an equation of the form [11, 23]

$$\ddot{\delta} + A(t)\dot{\delta} + B(t)\delta = 0. \quad (18)$$

Assuming that this equation remains valid when $\mathcal{R}$ goes infinite, the authors of [6] impose a scale-independent continuity condition on $\delta(t)$ across $t = 0$. This leads at $t > 0$ to a Bardeen potential with the mild scale dependence given by Eq. (14), and to a curvature perturbation which by the time of radiation domination has the same scale-dependence, which can agree with observation.

The failure of linear cosmological perturbation theory We will now show that the analysis just described is invalid, by establishing the following contradiction: an *assumption* of the analysis is that the Universe has practically uniform energy density and practically uniform (actually zero) spatial curvature, so that the equations of linear cosmological perturbation theory and in particular Eq. (18) can be derived, yet a *conclusion* of the analysis is that the above assumption is violated as the singularity is approached. To be more precise, we shall demonstrate that for every choice of slicing used to define the purportedly linear theory, either the density contrast or the curvature perturbation blows up as the singularity is approached (or both do), thus invalidating the linear theory with that choice.

The demonstration is very simple. We have seen already that on comoving slices, the curvature perturbation $\mathcal{R}$ blows up. Now consider a generic slicing, separated from the comoving one by time displacement $\delta t(\mathbf{x}, t)$. On such a slicing, the density contrast δ_s and the curvature perturbation $\mathcal{R}_s$ (defined as in Eq. (10)) are given by [11, 1]

$$\begin{aligned} \delta_s &= \delta - 3(1+w)|H|\delta t \\ \mathcal{R}_s &= \mathcal{R} + |H|\delta t. \end{aligned} \quad (19)$$

As the singularity is approached, $1 + w$ approaches the nonzero value 2, δ approaches a constant and $\mathcal{R}$ diverges. Therefore, no choice of δt can keep both δ_s and $\mathcal{R}_s$ finite as the singularity is approached.

One might think that the divergence of the curvature perturbation on comoving slices is by itself enough to invalidate the use of Eq. (18) since the relevant quantity δ has been defined as the density contrast on comoving slices. This definition obviously fails when the curvature of the slices diverges, and moreover the derivation of Eq. (18) uses Eqs. (15) and (17) which apparently require $\mathcal{R}$ to be small. However, within linear theory quantities defined in a given gauge can always be redefined in terms of quantities in any other gauge. (Recall that we gave an explicit example of this in the case of the Bardeen potential Φ .) Therefore, if linear theory were valid in just one gauge, it would be possible to redefine δ in terms of quantities in that gauge, and then Eq. (18) would be valid.

Conclusion It is difficult to generate the observed curvature perturbation in the pre-big-bang and ekpyrotic scenarios. It cannot be generated during collapse and there is no known mechanism for generating it at the bounce either. Subsequent generation by a suitable curvaton field is possible, but the curvaton must have a suitable non-canonical kinetic term and a suitable initial value. All this is likely to hold in any scenario where inflation is replaced by superluminal collapse, because it stems from the fact that the Hubble parameter is rapidly varying. By contrast the curvature perturbation is easily generated in the inflation scenario, either directly by the inflaton field or indirectly by a curvaton field.

Acknowledgement I thank David Wands for many helpful remarks about the evolution of the curvature perturbation. I have benefited also from correspondence with Jai-chan Hwang, Robert Brandenberger, Ruth Durrer and Gabrielle Veneziano.

References

- [1] D. H. Lyth and A. R. Liddle, *Cosmological Inflation and Large-Scale Structure*, Cambridge University Press (2000).
- [2] D. H. Lyth and D. Wands, hep-ph/0110002.
- [3] G. Veneziano, hep-th/0002094; R. Brustein, M. Gasperini, M. Giovannini, V. F. Mukhanov and G. Veneziano, Phys. Rev. D **51** (1995) 6744.
- [4] J. E. Lidsey, D. Wands and E. J. Copeland, Phys. Rep. **337** (2000) 343.
- [5] J. Khoury, B. A. Ovrut, P. J. Steinhardt and N. Turok, hep-th/0103239.
- [6] J. Khoury, B. A. Ovrut, P. J. Steinhardt and N. Turok, hep-th/0109050.
- [7] P. J. Steinhardt and N. Turok, hep-th/0110030 and hep-th/0111098.

- [8] D. H. Lyth, hep-ph/0106153,
- [9] D. H. Lyth and A. R. Liddle, Phys. Rep. **231** (1993) 1.
- [10] D. H. Lyth and A. Riotto, Phys. Rep. **314** (1998) 1.
- [11] J. M. Bardeen, Phys. Rev. D**22** (1980) 1882.
- [12] J. M. Bardeen, P. J. Steinhardt and M. S. Turner, Phys. Rev. D**28** (1983) 679.
- [13] H. Kodama and M. Sasaki, Prog. Theor. Phys. **78** (1984) 1 .
- [14] V. F. Mukhanov, Phys. Lett. **B218** (1989) 17.
- [15] M. Sasaki, Prog. Theor. Phys. **76** (1986) 1036 .
- [16] A. Buonanno, M. Lemoine and K. A. Olive, Phys. Rev. D**62** (2000) 083513.
- [17] K. Enqvist and M. S. Sloth, hep-ph/0109214.
- [18] R. Kallosh, L. Kofman and A. Linde, hep-th/0104073.
- [19] A. H. Guth and S.-Y. Pi, Phys. Rev. Lett. **49** (1982) 1110.
- [20] D. W. Olson, Phys. Rev. D**14** (1976) 327.
- [21] J. Khoury, B. A. Ovrut, N. Seiberg, P. J. Steinhardt and N. Turok, hep-th/0108187.
- [22] R. Brandenberger and F. Finelli, hep-th/0109001.
- [23] D. H. Lyth, Phys. Rev. D**31** (1985) 1792.